

Comparison of vitamins K₁, K₂ and K₃ effects on growth of rat glioma and human glioblastoma multiforme cells in vitro

Pinar OZTOPÇU, Selda KABADERE¹, Ayşe MERCANGÖZ, Ruhi UYAR¹

Osmangazi University Art and Sciences Faculty Department of Biology, and Medical Faculty Physiology Department¹ Eskişehir, Türkiye

Abstract

Glioblastoma multiforme is characterized as highly invasive and rapidly growing astrocytomas, and scientists have sought for efficient treatment against malignant gliomas for a long time. Therefore, we compared the response of rat glioma (C6) and glioblastoma multiforme cells derived from two patients to vitamins K₁, K₂ and K₃. The cells were exposed to 100, 250, 500, 750 and 1000 µM of vitamins K₁ and K₂, and 1, 10, 25, 50, 75 and 100 µM of vitamin K₃ for 24 hours in an incubator atmosphere of 5% CO₂, 37°C and 100% humidity. Cell viability was estimated by MTT assay. Vitamin K₁ showed no growth effect on all the glioma cells examined. Vitamin K₂ did not cause any change in number of C6, however induced growth inhibition in a dose-dependent manner on glioblastoma multiforme. The IC₅₀ values of vitamin K₂ were 960 µM and 970 µM for glioblastoma multiforme, respectively. Vitamin K₃ had also growth inhibitory effect in a dose-dependent manner on both C6 and glioblastoma multiforme. The IC₅₀ values were 41 µM, 24 µM and 23 µM for vitamin K₃, respectively. We concluded that vitamin K₃ is more effective than vitamin K₂ for inhibition of cancer cell growth, and might have an alternative value as an anti-cancer drug against glioblastoma multiforme.

Key words: Glioma ; K vitamins ; cell proliferation ; MTT assay ; in vitro.

1. Introduction

Glioblastoma multiforme is characterized as rapidly growing and highly invasive astrocytomas (Louis and Cavenee, 1997). It is more common in older adults and affects more men than women. These tumor types can be treated with surgery, chemotherapy and radiation therapy. Since cells of the tumor stray into other areas of the brain, surgery alone rarely controls the glioblastoma multiforme (Mikkelsen, 1998). Several classes of drugs are used for the chemotherapy of this type of tumors such as alkylating agents, nitrosoureas or polyamine inhibitors, however clinical resistance to chemotherapy and side effects have been major problems in treatment (Salford, 2002). The mean survival time after diagnosis for patients with glioblastoma multiforme is less than a year in spite

of surgery, radiotherapy and chemotherapy (Kimmel, 1987). For a long time scientists have sought for efficient treatment against malignant gliomas.

Vitamin K exists in at least three different forms : vitamin K₁ (phylloquinone ; 2-methyl-3-phytyl-1,4 naphthoquinone) mostly found in green leafy vegetables, vitamin K₂ (menaquinone ; 2-methyl-3-(all-trans-difarnesyl)-1,4 naphthoquinone) synthesized by the intestinal flora, and vitamin K₃ (menadione ; 2-methyl-1,4 naphthoquinone) a synthetic derivative of vitamin K₁ and K₂ (Suttie, 1995).

Previous studies demonstrated that vitamin K₃ inhibits cell growth of various rodent and many types of malignant human tumor cells in vivo and in vitro (Nishikawa *et al.*, 1995 ; Ngo *et al.*, 1991). Vitamin K₁ and K₂ also have been shown to have growth inhibitory effects in vitro, but these effects are much weaker than those of vitamin K₃ (Prasad *et al.*, 1981 ; Wu *et al.*, 1993). Beside a broad spectrum of antitumor activity vitamin K₃ was shown to enhance the cytotoxic effect of some clinically useful anticancer agents, including action against multidrug resistant human cancer cell lines, and it was thought to have less serious toxic effects (Nutter *et al.*, 1991 ; Wang *et al.*, 1995). We noticed that there are insufficient studies related to the effect of K vitamins on growth of glioma cells in vitro. In addition there isn't any study comparing the antiproliferative effect of K vitamins on both rat glioma cell line and human glioblastoma multiforme cells. Therefore, we have attempted to compare the possible inhibitory effect of vitamin K₁, K₂ and K₃ based on MTT assay using both C6 and low passage human glioblastoma multiforme cells obtained from two different patients tumor tissue in culture.

2. Methods and materials

2.1. REAGENTS AND CELL CULTURE

All chemicals were purchased from Sigma Chemical Co. (St. Louis, MO). We used C6 and cultured low passage human glioblastoma

multiforme which were derived from primary cultures of two patients' tumor tissue as described previously by Durmaz *et al.* (Durmaz *et al.*, 1999).

All the cells were maintained in 75 cm² flasks, and incubated in 1:1 mixture of Dulbecco's Modified Eagle's Medium and F-12 supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin solution at 37°C in a humidified atmosphere of 5% CO₂ and 95% air (Unterman *et al.*, 1991). The medium was changed 2-3 times within a week until the cultures approached confluence. For experimentation, monolayer cultures were washed with Hank's balanced salt solution (without calcium and magnesium) and then incubated in the presence of 0,25% 3 ml trypsin-EDTA solution for 10-15 minutes at 37°C. After the cells were dispersed, trypsin activity was inhibited by adding growth medium and the cells were centrifuged at 1000 rpm for 5 minutes at 4°C, and then counted with a Coulter counter. Cell viability was accessed by trypan blue dye exclusion and found to be higher than 98%. Vitamin K₁ and K₂ were dissolved in absolute ethanol at 1000 µM concentration. Further dilutions were made at ratio of 1:10 in F-12 and the maximum concentration of ethanol was adjusted to be 0,01%. This amount was added to the culture medium together with the highest concentrations of vitamins K₁ and K₂ (1000 µM), and found to have no effect on cell viability when used alone (data not given). Water soluble form of vitamin K₃ (menadione sodium bisulfate) was dissolved in F-12 medium. For experimental protocol, the cells at exponential growth phase were seeded in 3 x 10⁴ cells/well in 96 well microtiter plates, 10 wells for control (included growth medium only) and 10 wells for each tested K vitamins doses, and incubated for 24 hours. Vitamin K₁ and K₂ at the concentrations of 100, 250, 500, 750, 1000 µM; vitamin K₃ at the doses of 1, 10, 25, 50, 75, 100 µM were added to the growth medium for 24 hours. All the test compounds were prepared immediately prior to use and protected from light.

2.2. MTT COLORIMETRIC ASSAY

After 24 hours, drug cytotoxicity screening was determined by using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) colorimetric assay originally described by Mossmann and modified by Alley *et al.* (Mossmann, 1983; Alley *et al.*, 1988). MTT was prepared freshly before each test as 5 mg/ml in phosphate buffered saline and filtered to sterilize with 0,22 µm filter. Then, 25 µl MTT solution was added to the each well and incubated for 4 hours at 37°C. The MTT solution was converted into blue formazan by mitochondrial dehydrogenase activity of viable cells (Alley *et al.*, 1988). Therefore, the amount of formazan produced is proportional to the number of living cells (Abe and Matsuki, 2000). After the




FIG. 1. — Action of vitamin K₁ on C6 cells, low passage human glioblastoma multiforme cells from the first patient (GBM-1) and human glioblastoma multiforme cells from the second patient (GBM-2).

medium was removed from the wells, 100 µl dimethyl sulphoxide was added to each well and the crystal formazan particles produced in viable cells was dissolved for 5 minutes at room temperature with a shaker. The absorbance of formazan dye was read at 550 nm using a microplate reader (Bio-Tek Instruments), and cell survival percentages were calculated according to the following formula; absorbance of treated cells in each well x 100 / mean absorbance of control cells (Durmaz *et al.*, 1999; Loveland *et al.*, 1992). The dose response curves were calculated for three vitamin K forms at the above-mentioned concentrations and expressed as the mean percent fraction of control ± SEM. Fifty % growth inhibitory concentration (IC₅₀) was determined by plotting the logarithm of the drug concentration against the mean percentage of living cells at the each dose of K vitamins. All statistical analyses were performed using one-way analysis of variance (ANOVA) and followed by Tukey's multiple comparison tests. A p value less than 0.05 was considered to be significant. Each experiment was repeated at least three times.

3. Results

When compared to the control, vitamin K₁ did not show any activity on C6 and the low passage human glioma cells proliferation (Fig. 1). K₂ also had no effect on C6 cell proliferation, however it inhibited cell survival in a dose-dependent manner in the human glioma cells. Vitamin K₂ at the doses of 100, 250 and 500 µM had no effect on the survival of the human glioma cells, but inhibitory effect was observed at 750 µM and reached down to 56% reduction at 1000 µM (p < 0.001). IC₅₀ of vitamin K₂ for the human glioma cells were esti-




FIG. 2. — Effects of vitamin K₂ on C6, GBM-1 and GBM-2 cells. A means $p < 0.001$ when compared to the control.




FIG. 3. — Cytotoxic effects of vitamin K₃ on C6, GBM-1 and GBM-2 cells.

ated as 960 µM and 970 µM, respectively (Fig. 2).

Vitamin K₃ at the concentrations of 1, 10 and 25 µM (Fig. 3) did not influence C6 cell viability when compared to the control, but decreased the cell number starting from 50 µM and reached to almost 100% reduction at 100 µM ($P < 0.001$). IC₅₀ of vitamin K₃ for C6 cells was calculated as 41 µM. In contrast, vitamin K₃ reduced the cell number starting from 10 µM in the human glioma cells (Fig. 3). The decline in cell viability reached down to 90% at 50 µM ($p < 0.001$). IC₅₀ of vitamin K₃ for the human glioma cells was calculated as 24 µM and 23 µM, respectively. When compared to each other, there was no difference in terms of response of two different sources of human glioma cells to three different K vitamins at every dose studied.

4. Discussion

Vitamins of group K have pronounced cytotoxic properties leading to their application in the therapy of various neoplasms. Wu *et al.* compared the growth inhibitory effects of vitamins K₁, K₂ and K₃

against on five hepatoma cell lines, nasopharyngeal carcinoma, leukemia, oral epidermoid carcinoma and breast carcinoma cells. For all these cells, the ID₅₀ values of vitamin K₁ ranged from 6 to 9 mM, and the ID₅₀ of vitamin K₂ ranged from 1 to 2 mM. However, the respective ID₅₀ values of vitamin K₃ for these nine different cells were between 15 and 42 µM (Wu *et al.*, 1993). Nishikawa *et al.* reported that K vitamins inhibit growth of human hepatoma cells and 50% inhibitory doses were 1000 µM for vitamin K₁, 112 µM for K₂ and 10 µM for K₃ (Nishikawa *et al.*, 1995). Markovits *et al.* also evaluated the action of K vitamins on another human hepatoma cells and found IC₅₀ values approximately 500, 150 and 5 µM for vitamin K₁, K₂ and K₃, respectively (Markovits *et al.*, 2003). Sata *et al.* found that menadione at 1 and 2 µM doses, and time dependently inhibited proliferation of rat pancreatic acinar cell line (Sata *et al.*, 1997). Our results demonstrate that vitamin K₁ showed no growth effect on both C6 and human glioblastoma multiforme cells. Vitamin K₂ also did not show any growth inhibitory effect on C6, however induced growth inhibition in a dose-dependent manner on human glioma cells. Vitamin K₃ had antiproliferative effect on both C6 and human glioma cells. IC₅₀ of vitamin K₃ were estimated as 41 µM for C6 cells, 24 and 23 µM for human glioblastoma multiforme cells. All these previous and our results demonstrate that the growth inhibitory effect of vitamin K₃ is stronger as much as 1000-fold than vitamin K₁ or K₂ on different cells in vitro.

Generation of reactive oxygen species (ROS) by redox cycling is generally accepted as a mechanism of vitamin K₃ cytotoxicity (Buckman *et al.*, 1993). Recent reports indicated that especially astrocytes appear to be more resistant to ROS and play an important role in reducing ROS level in brain to protect neurons. Desagher *et al.* showed that the protective effect of astrocytes depend on the astrocytes/neurons ratio, and estimated that only one astrocytes has the capacity to significantly protect 20 neurons against the toxicity induced by a 30 min application of 100 µM hydrogen peroxide on mouse neuronal cultures (Desagher *et al.*, 1996). Moreover, Abe and Saito examined the effect of 100 µM K₃ in cultured rat cortical astrocytes and found apparent morphological changes 2 hours later, and all cells were killed in 12 hours (Abe and Saito, 1996). This concentration can be very high for treatment or combination. Our results suggested that vitamin K₃ inhibits cell growth at low concentrations on glioma cells. The amount of ROS formed by low levels of vitamin K₃ may not be enough to injure neurons. Meanwhile Sun *et al.* results suggested that when antioxidant enzymes were added prior to addition of vitamin K₃, toxicity was markedly attenuated (Sun *et al.*, 2000).

Both in vitro and in vivo studies for over half a century showed a synergistic or additive effect

when vitamin K₃ was combined with chemotherapeutic agents of about 20 different drugs on about 10 different tumors or cell lines (Lamson and Plaza, 2003). It is also concluded that vitamin K₃ has antitumoral activity on multi-drug resistant human cancer cell lines and thought to have less serious toxic side effects (Nutter *et al.*, 1991 ; Wang *et al.*, 1995). All these findings suggest that vitamin K₃ may be useful in combination treatment with antitumor drugs and some types of drug-resistant brain tumors. Although diffusion of chemotherapeutic drugs within the blood-brain barrier is fairly poor, vitamin K₃ has increased the permeability of rat brain capillary endothelial cells in an in vitro model of the blood brain barrier (Lagrange *et al.*, 1999).

In conclusion our results indicated that vitamin K₃ has potential antiproliferative effect on both C6 and the low passage human glioma cells. When the two cell types compared, antiproliferative effect of vitamin K₃ was stronger on the human glioma cells than on C6 cells. All these findings suggest that vitamin K₃ might be a good candidate in the clinical treatment of glioblastoma multiforme. For this reason, a study about the possible effects of vitamin K₃ on normal rat tissues is underway in our laboratory.

List of abbreviations

MTT : 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide.

EDTA : Ethylenediaminetetraacetic acid.

IC₅₀ : Fifty % growth inhibitory concentration.

ANOVA : Analysis of variance.

ID₅₀ : Fifty % growth inhibitory dose.

ROS : Reactive oxygen species.

Acknowledge

This study was supported by a grant (2001/6) from the Scientific Research Projects Commission of Osmangazi University, Eskişehir, Turkey.

REFERENCES

- ABE K., MATSUKI N. Measurement of cellular 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) reduction activity and lactate dehydrogenase release using MTT. *Neurosci. Res.*, 2000, **38** : 325-29.
- ABE K., SAITO H. Menadione toxicity in cultured rat cortical astrocytes. *Jpn. J. Pharmacol.*, 1996, **72** : 299-304.
- ALLEY M. C., SCUDIERO D. A., MONKS A., HURSEY M. L., CZERWINSKI M. J. Feasibility of drug screening with panels of human tumor cell lines using a microculture tetrazolium assay. *Cancer Res.*, 1988, **48** : 589-601.
- BUCKMAN T. D., SUTPHIN M. S., MITROVIC B. Oxidative stress in a clonal cell line of neuronal origin : effects of antioxidant enzyme modulation. *J. Neurochem.*, 1993, **60** (6) 2046-58.
- DESAGHER S., GLOWINSKI J., PREMONT J. Astrocytes protect neurons from hydrogen peroxide toxicity. *J. Neurosci.*, 1996, **16** (8) : 2553-62.
- DURMAZ R., DELIORMAN S., UYAR R., ISIKSOY S., EROL K. The effects of anticancer drugs in combination with nimodipine and verapamil on cultured cells of glioblastoma multiforme. *Clin. Neurol. Neurosurg.*, 1999, **101** : 238-44.
- KIMMEL D. W., SHAPIRO J. R., SHAPIRO W. R. In vitro drug sensitivity testing in the human gliomas. *J. Neurosurg.*, 1987, **66** : 161-71.
- LAGRANGE P., ROMERO I. A., MINN A., REVEST P. A. Transendothelial permeability changes induced by free radicals in an in vitro model of the blood-brain barrier. *Free Rad. Biol. Med.*, 1999, **27** : 667-72.
- LAMSON D. W., PLAZA S. M. The anticancer effects of vitamin K. *Altern. Med. Rev.*, 2003, **8** (3) : 303-18.
- LOUIS D. N., CAVENEE W. K. Neoplasms of the central nervous system. In : DEVITA V.T., HELLMAN S., and ROSENBERG S. A. (eds.). *Cancer : principle and practice of oncology*, 5th ed. Philadelphia : Lippincott-Raven Publishers, 1997, 2013-17.
- LOVELAND B. E., JOHNS T. G., MACKAY I. R., VAILLANT F., WANG Z.-X., HERTZOG P. J. Validation of the MTT dye assay for enumeration of cells in proliferative and antiproliferative assays. *Biochem. Int.*, 1992, **27** (3) : 501-10.
- MARKOVITS J., WANG Z., CARR B. I., SUN T. P., MINTZ P., BRET M. L., WU C.-W., WU F. Y.-H. Differential effects of two growth inhibitory K vitamin analogs on cell cycle regulating proteins in human hepatoma cells. *Life Sci.*, 2003, **72** : 2769-84.
- MIKKELSEN T. Cytostatic agents in the management of malignant gliomas. *Cancer Control : JMCC*, 1998, **5** (2) : 150-62.
- MOSMANN T. Rapid colorimetric assay for cellular growth and survival : Application to proliferation and cytotoxicity assay. *J. Immunol. Methods*, 1983, **65** : 55-63.
- NGO E. O., SUN T. P., CHANG J. Y., WANG C. C., CHI K. H., CHENG A. L., NUTTER L. M. Menadione-induced DNA damage in a human tumor cell line. *Biochem. Pharmacol.*, 1991, **42** (10) : 1961-68.
- NISHIKAWA Y., CARR B. I., WANG M., KAR S., FINN F., DOWD P., ZHENG Z., KERNS J., NAGANATHAN S. Growth inhibition of hepatoma cells induced by vitamin K and its analogs. *J. Biol. Chem.*, 1995, **270** (47) : 28304-10.
- NUTTER L. M., CHENG A. L., HUNG H. L., HSIEH R. K., NGO E. O., LIU T.-W. Menadione : spectrum of anticancer activity and effects on nucleotide metabolism in human neoplastic cell lines. *Biochem. Pharm.*, 1991, **41** : 1283-92.
- PRASAD K. N., PRASAD J. E., SAKAMOTO A. Vitamin K₃ (menadione) inhibits the growth of mammalian tumor cells in culture. *Life Sci.*, 1981, **13** : 1387-92.
- SALFORD L. G., SIESJÖ P., SKAGERBERG G., PERSSON B. R. R., LARSSON E. M. Search for effective therapy against glioblastoma multiforme-clinical immunisation with autologous glioma cells transduced with the human interferon- γ

- gene. *International Congress Series*, 2002, **1247** : 211-20.
- SATA N., STUMPE K. H., HAN B., HAUSSINGER D., NIEDERAU C. Menadione induces both necrosis and apoptosis in rat pancreatic acinar AR4-2J cells. *Free Rad. Biol. Med.*, 1997, **23** (6) : 844-50.
- SUN L. K., YOSHII Y., MIYAGI K. Cytotoxic effect through Fas/APO-1 expression due to vitamin K in human glioma cells. *J. Neurooncol.*, 2000, **47** : 31-8.
- SUTTIE J. W. The importance of menaquinones in human nutrition. *Annu. Rev. Nutr.*, 1995, **15** : 399-417.
- UNTERMAN T. G., GLICK R. P., WAITES G. T. Production of insulin-like growth factor-binding proteins by human central nervous system tumors. *Cancer Res.*, 1991, **51** : 3030-36.
- WANG Z., WANG M., FINN F., CARR B. I. The growth inhibitory effects of vitamin K and their actions on gene expression. *Hepatology*, 1995, **22** (3) : 876-82.
- WU F. Y.-H., LIAO W.-C., CHANG H.-M. Comparison of antitumor activity of vitamins K₁, K₂ and K₃ on human tumor cells by two (MTT and SRB) cell viability assays. *Life Sci.*, 1993, **52** (22) : 1797-1804.

R. UYAR,
Osmangazi University,
Medical Faculty,
Physiology Department,
26480 Eskişehir (Türkiye).
E-mail : ruyar@ogu.edu.tr.